

Rodent Prey of the Barn Owl *Tyto alba* and Short-Eared Owl *Asio flammeus* During Winter in Agricultural Lands in Southern Chile

Author(s): Ricardo A. Figueroa R., Jaime R. Rau, Sonia Mayorga, David R. Martínez, E. Soraya Corales S., Andrés Mansilla, & Rodolfo Figueroa M.

Source: *Wildlife Biology*, 15(2):129-136.

Published By: Nordic Board for Wildlife Research

DOI: <http://dx.doi.org/10.2981/08-005>

URL: <http://www.bioone.org/doi/full/10.2981/08-005>

BioOne (www.bioone.org) is a nonprofit, online aggregation of core research in the biological, ecological, and environmental sciences. BioOne provides a sustainable online platform for over 170 journals and books published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

Rodent prey of the barn owl *Tyto alba* and short-eared owl *Asio flammeus* during winter in agricultural lands in southern Chile

Ricardo A. Figueroa R., Jaime R. Rau, Sonia Mayorga, David R. Martínez, E. Soraya Corales S., Andrés Mansilla & Rodolfo Figueroa M.

We compared the consumption of rodent prey by barn owl *Tyto alba* and short-eared owl *Asio flammeus* during winter in agricultural areas in southern Chile. Diets were studied on the basis of pellets collected during the winters of 1986, 1987 and 1996. Both owl species consumed a large number of different rodent prey (8-9 species), but they preyed more often on the olivaceous field mouse *Abrothrix olivaceus* (56% and 52% of all individual prey, respectively) and long-tailed pygmy rice rat *Oligoryzomys longicandatus* (12.6% and 18.6%, respectively). The diet of the two owl species largely overlapped (95%). The diet diversity of barn owls and short-eared owls was not statistically different and accordingly they showed a similar evenness in diet. The geometric mean weight of rodent prey (GMWP) for short-eared owls (33 ± 1.36 g) was significantly greater than that of barn owls (28.3 ± 1.81 g). No significant difference was found between the proportion of native and introduced rodent prey consumed by barn and short-eared owls. The high diet similarity between both owl species could be a result of convergencies in hunting modes and activity time, similar body mass, or homogeneous rodent prey distribution and abundance in the agricultural areas we studied. The higher GMWP for the short-eared owl was probably caused by the fact that it preyed on the largest rodent prey in our study sites, the Norwegian rat *Rattus norvegicus*. According to our rodent trapping in the field, barn and short-eared owls appear to be opportunistic rather than selective predators in agricultural areas in southern Chile.

Key words: agricultural lands, *Asio flammeus*, barn owl, diet overlap, rodent prey, short-eared owl, *Tyto alba*

Ricardo A. Figueroa R., Instituto de Zoología, Escuela de Graduados, Facultad de Ciencias, Universidad Austral de Chile, Casilla 567, Valdivia, Chile - e-mail: ra_figueroa_rojas@yahoo.com

Jaime R. Rau & Sonia Mayorga, Laboratorio de Ecología, Depto Ciencias Básicas & Programa IBAM, Universidad de Los Lagos, Osorno, Chile - e-mail address: arau@telsur.cl (Jaime R. Rau)

David R. Martínez, CEFORA, Universidad de Los Lagos, Osorno, Chile - e-mail: dmartin@ulagos.cl

E. Soraya Corales S., Estudios para la Conservación y Manejo de la Vida Silvestre Consultores, Saavedra 1199, Victoria, Chile - e-mail: sorayacoraless@yahoo.com

Andrés Mansilla, Departamento de Ciencias y Recursos Naturales, Facultad de Ciencias, Universidad de Magallanes, Punta Arenas, Chile - e-mail: andres.mansilla@umag.cl

Rodolfo Figueroa M., Instituto de Ecología y Evolución, Facultad de Ciencias, Universidad Austral de Chile, Casilla 567, Valdivia, Chile - e-mail: rfigueroa@uach.cl

Corresponding author: Ricardo A. Figueroa R.

Received 18 January 2008, accepted 19 November 2008

Associate Editor: Klaus Hackländer

Dietary studies of raptors have ecological importance, because they may help understand prey distribution, abundance, behaviour and vulnerability (Fulk 1976, Marti 1987, Torre et al. 2004), energetic requirements (Bozinovic & Medel 1988) or trophic

relationships between sympatric species and raptor assemblage structures (Herrera & Hiraldo 1976, Jaksic & Braker 1983, Jaksic 1985). Factors that could influence interspecific trophic relationship and potential mechanisms of food partitioning among

coexisting raptor species are the abundance, habitat selection and body mass of prey, and feeding strategies, body size, hunting range, territoriality and activity time of raptors (Jaksic 1989, Gerstell & Bednarz 1999, Leveau et al. 2004, García & Arroyo 2005). Generally, coexisting raptor species show a low food-niche overlap when prey are in scarce supply which is assumed as the presence of competition by some authors (e.g. Lack 1946, Steenhof & Kochert 1985, García & Arroyo 2005). On the other hand, a high food-niche overlap may occur when prey resources increase, which is generally interpreted as opportunistic convergence on abundant resources (Lack 1946, Walk 1998, Leveau et al. 2004). The territoriality or agonistic behaviour may become an important factor in food-niche segregation (Jaksic 1985), but this could vanish with high food supply. The existence of different hunting modes or feeding strategies among raptors could also lead to food-niche segregation (Jaksic & Carothers 1985). If this is true, similar hunting modes among raptor birds would lead to food-niche convergence. Food partitioning by prey size might occur between species that differ in body size, with larger prey expected to be preferred with increasing body size of the raptor (Jaksic & Braker 1983, Marti 1992, Bellocq & Kravetz 1994, García & Arroyo 2005). In this case, it would be expected that raptors of similar body size take the same average prey size. Although time of activity has been demonstrated as a factor potentially involved in resource partitioning (Marti & Kochert 1995), it does not appear to be sufficiently adequate to separate food-niches of birds of prey (Jaksic 1982). Many raptors hunt during the crepuscule when both diurnal and nocturnal prey are active, reducing differences in resource partitioning.

Although barn owl *Tyto alba* and short-eared owl *Asio flammeus* are sympatric over most of their range in Chile, their diets have never been compared. The few published works on the diet of both owl species in agricultural lands of southern Chile indicate that they mainly prey upon small rodents (Rau et al. 1985, 1992, Martínez et al. 1998). Since both owl species have similar body mass (350-400 g) and hunting modes, search for prey in the same habitats, and may overlap in activity time, they should have access to similar prey species. In addition, when there is an ample supply of prey, barn owls and short-eared owls should have high dietary similarities. Here, we examine these hypotheses by evaluating rodent consumption by coexistent barn owls and short-eared owls in agricultural lands of south-

ern Chile during the period of highest seasonal rodent abundance (winter).

Study area

Our study was conducted in four agricultural areas of the Osorno province, Central Valley in southern Chile. Localities were Quirislahuén, Pilauco, Chuyaca and Chahuilco. The first three localities are very close to Osorno city (3-4 km) and the latter is located 15 km south of Osorno (40°44'S, 73°09'W). In general, the landscape is relatively homogeneous with almost all agricultural areas presenting a similar topography, vegetation and land management (Martínez et al. 1998). Topography is characterised by undulating terrains and extensive flat areas. Vegetation is composed of mainly pasturelands for livestock (e.g. cattle), croplands (e.g. wheat *Triticum aestivum*, oat *Avena sativa* and potatoes *Solanum tuberosum*), and fruit and legume gardens associated with the dispersed human settlements. Scattered small (1-4 ha) abandoned pastureland with dispersed isolated and old native trees (*Nothofagus obliqua* and *Laurelia sempervirens*) are frequent between the currently used fields. Webs of fallow vegetation strips continuously cover boundaries between agricultural fields and road borders, and some small and altered patches of the original native *Nothofagus* forest border rivers and streams. The flat terrain contains sedge-rush *Carex-Juncus* marshes (seasonal or permanent). The climate is moist-temperate with a Mediterranean influence, characterised by heavy rainfall (1500-3000 mm annually), wet, frost-free winters and short, dry summers (di Castri & Hajek 1976). Annual mean temperature is 12°C.

Material and methods

The barn owls' pellets were collected from Pilauco (May-July 1986, N=26), Quirislahuén (May-June 1987, N=42), and Chahuilco (May-July 1996, N=160). Samples were taken either under old native trees where barn owls perched or in tree cavities where they roosted. We made sure that pellets were from barn owls by direct observation of birds and permanent use of site collection. Information on short-eared owls was taken from previous works (Rau et al. 1992, Martínez et al. 1998), in which short-eared owl pellets were collected from

Quirislahuén (May-June 1986, N=46), Chuyaca (May-August 1995, N=147), and Chahuilco (May-August 1995, N=64). Samples were taken on open grounds in abandoned pastures or dried marshes, and under trees, fences and stumps used as perches. Confirmation that pellets belonged to short-eared owls was made by flushing or by using indirect signs on perches (e.g. feathers and creamy fecal waste; Holt et al. 1987). As wintering barn and short-eared owls largely depend on small mammals, other occasional prey (e.g. birds and insects) were discarded from the analysis because of low biomass contribution. We only analysed winter rodent consumption, because it was not possible to obtain pellets year-round for all localities or because samples were very small.

Rodent prey were identified and quantified on the basis of skull or dentary pairs following keys in Reise (1973) and Pearson (1995). For hair remains, we used reference collections and quantified these prey assuming the smallest possible number of individuals (e.g. hair of given a species was deemed as representing only one individual). We identified prey items to the finest possible taxonomic category in all cases. Biomass contribution of each prey species was estimated by multiplying the number of individuals in the pellets by mean body mass of each species (Marti 1987). Masses of rodent prey were taken from the unpublished database of the Ecology and Evolution Institute of the Austral University of Chile in Valdivia, Chile, and Agricultural and Livestock Service in Aysén, Chile. Body masses of rodent species not available in databases were taken from the literature. We assumed that masses of unidentified prey were similar to the mean mass of the most closely related identified taxon.

Diet of owls were compared utilising food-niche estimators (Marti 1987). The diet diversity (food-niche breadth) was estimated by use of the Shannon-Wiener Index (H' ; Krebs 1989) and statistically compared by the Hutcheson t-test (Zar 1996). Complementarily, we evaluated how similar the frequencies of individual prey in diet were by using the index of evenness based on the Shannon-Wiener function: $J' = H' / \log S$; where S is the number of species in the sample (Krebs 1989). The J' index ranges from 0 (no prey item is equally represented) to 1 (all prey items are equally represented). The diet similarity (food-niche overlap) was estimated by using the Horn index (R_o ; Krebs 1989). This index ranges from 0 (no similarity) to 1 (complete similarity). The geometric mean weight of prey (GMWP) was calculated fol-

lowing Marti (1987): $GMWP = \text{antilog}(\sum n_i \log w_i / \sum n_i)$, where n_i is the number of individuals of the i th species, and w_i is the mean weight. Although some authors have suggested that geometric means tend to underestimate mean weight of prey (Walker et al. 2007), we preferred using GMWP, because it made comparisons with previous studies on owls diet possible. In addition, differences in consumption of introduced (Murinae) versus native (Sigmodontinae) rodents were evaluated with χ^2 -tests using contingency tables (Fowler & Cohen 1986).

Except for Chuyaca, we were unable to obtain concurrent data on the field abundance of rodent prey in our study localities. In Chuyaca, we evaluated the relative abundance of rodents by live-trapping transects using medium Sherman traps in abandoned pastures located inside the owls' hunting area (see Martínez et al. 1998 for details). Because this trapping was locally restricted and only a small fraction of owls's prey was captured, we combined results from Chuyaca with data collected from three other vegetationally similar agricultural areas by using the same trapping method to obtain a coarse estimate of the rodent field abundance (Fig. 1). Trapping was made either in pasture, marshes or




Figure 1. Comparison of the proportion of rodent prey in the winter diet of barn and short-eared owls versus the winter rodent abundance in the field estimated by live-trapping transects in agricultural areas of southern Chile. Alon = *Abrothrix longipilis*, Aoli = *A. olivaceus*, Gval = *Geoxus valdivianus*, Mmus = *Mus musculus*, Olon = *Oligoryzomys longicaudatus*, Lmic = *Loxodontomys micropus*, Rnor = *Rattus norvegicus*, Rrat = *R. rattus* and Pdar = *Phyllotis darwini*.

forest edges to better represent the prey availability on the owls' hunting areas. However, because most abundance estimates were obtained five or six years after pellet collections or from areas where owls did not hunt, no statistical comparisons were made.

Results

In all localities, the barn owls mostly preyed upon the olivaceous field mouse *Abrothrix olivaceus* and secondarily upon the southern big-eared mouse *Loxodontomys micropus* or the long-tailed pygmy rice rat *Oligoryzomys longicaudatus* (Table 1). The remaining rodent prey account for less than 8% by number for all localities (see Table 1). The olivaceous field mouse also accounted for the higher contribution of biomass, followed by the southern big-eared mouse and introduced rats (*Rattus* spp., see Table 1). The barn owl diet was highly similar among localities with $R_o=0.90$ for Pilauco and Quirislahuén, $R_o=0.89$ for Pilauco and Chahuilco, and $R_o=0.89$ for Quirislahuén and Chahuilco. The diet of short-eared owls was also highly similar among localities (symmetrical niche overlap=91.8%; Martínez et al. 1998). Thus, we combined results from all localities for comparisons between both owl species.

Overall, barn owls preyed on at least eight rodent species with olivaceous field mouse being the most important prey by both number and biomass (Table 2). Taking into account both number and biomass, the southern big-eared mouse and long-tailed pygmy rice rat were the secondary prey of barn owls (see

Table 2). Although introduced rats were less consumed, their biomass contribution was similar to that of southern big-eared mice (see Table 2). Short-eared owls preyed upon nine rodent species with olivaceous field mouse being the most important by number and second by biomass (see Table 2). In spite of the fact that the number contribution of the Norwegian rat *Rattus norvegicus* was low, it accounted for the highest biomass contribution (see Table 2). As with barn owl, the southern big-eared mouse and long-tailed pygmy rice rat were the secondary prey both in terms of number and biomass (see Table 2). The diets of both owl species overlapped considerably ($R_o=0.95$). The diet diversity of barn owl and short-eared owl was not statistically different ($t_{200}=0.94$, $P>0.05$), and accordingly, they showed a similar evenness (see Table 2). The GMWP for short-eared owls (33 ± 1.36 g) was significantly greater than that of barn owls (28.3 ± 1.81 ; $t_{745}=-4.32$, $P<0.01$). No significant difference was found between the proportion of native and introduced rodents in barn owl diet from different localities ($\chi^2=0.3$, $P>0.05$), or for barn versus short-eared owls' diet ($\chi^2=1.1$, $P>0.05$).

A total of 211 individuals, including seven different rodent species, were captured in 1664 trap nights. The most captured species were the olivaceous field mouse and long-tailed pygmy rice rat which accounted for 47% and 36% of all individual rodents, respectively. The southern big-eared mouse was the least captured species (<1% of all individual rodents). In general, the ranking of prey species in the diet of barn and short-eared owls reflected the

Table 1. Winter diet of the barn owl *Tyto alba* in three agrosystems in southern Chile. Rodent mass is given as mean \pm SE. N%= number percentage and B%= biomass percentage.

Species	Mass (g)	Quirislahuén		Pilauco		Chahuilco	
		N%	B%	N%	B%	N%	B%
<i>Abrothrix longipilis</i>	33.4 \pm 0.26	1.1	0.9	7.3	6.8	0.3	0.3
<i>Abrothrix olivaceus</i>	24.8 \pm 0.28	43.2	24.6	60.0	41.6	58.5	39.2
<i>Geoxus valdivianus</i>	29.1 \pm 2.5	0.0	0.0	0.0	0.0	0.6	0.5
<i>Loxodontomys micropus</i>	50.7 \pm 1.4	16.0	18.5	10.9	15.5	7.1	9.8
<i>Oligoryzomys longicaudatus</i>	27.8 \pm 0.25	4.5	3.0	7.3	5.6	15.8	11.8
<i>Phyllotis darwini</i>	66 ^a	1.1	1.7	0.0	0.0	0.0	0.0
<i>Rattus rattus</i>	107.5 \pm 16.9	4.5	11.2	5.4	16.4	0.0	0.0
<i>Rattus</i> spp.	150 ^b	1.1	3.9	0.0	0.0	5.0	20.1
<i>Mus musculus</i>	14.6 \pm 1.37	0.0	0.0	0.0	0.0	0.6	0.2
Unidentified rodents	55.5 ^c	28.5	36.2	9.0	14.1	12.1	18.1
Total prey items		88		55		323	
Total biomass (g)		3830.3		1967.4		11956.4	
Total pellets		42		26		160	

^a Mass obtained from Jaksic et al. 1986.

^b Mean mass of the two *Rattus* species inhabiting study areas (*R. rattus* and *R. norvegicus*).

^c Mean mass of all identified rodent prey.

Table 2. Rodent prey of the barn owl and short-eared owl during winter in agrosystems in southern Chile. Rodent mass is given as mean \pm SE. N% = number percentage and B% = biomass percentage.

Prey species	Mass (g)	Barn owl		Short-eared owl	
		N%	B%	N%	B%
<i>Abrothrix longipilis</i>	33.4 \pm 0.26	1.3	1.1	2.0	1.5
<i>Abrothrix olivaceus</i>	24.8 \pm 0.28	56.0	36.3	51.7	28.8
<i>Geoxus valdivianus</i>	29.1 \pm 2.5	0.4	0.3	1.0	0.6
<i>Loxodontomys micropus</i>	50.7 \pm 1.4	9.2	12.3	8.7	9.9
<i>Oligoryzomys longicaudatus</i>	27.8 \pm 0.25	12.6	9.2	18.6	11.6
<i>Phyllotis darwini</i>	66 ^a	0.2	0.4	0.3	0.4
<i>Rattus norvegicus</i>	201 ^b	0.0	0.0	7.4	33.6
<i>Rattus rattus</i>	107.5 \pm 16.9	1.5	4.2	0.7	1.8
<i>Rattus</i> spp.	150 ^c	3.6	14.4	0.0	0.0
<i>Mus musculus</i>	14.6 \pm 1.37	0.4	0.2	0.2	0.1
Unidentified rodents	41.7 ^d	14.8	21.6	9.4	11.7
Total prey items		466		404	
Total biomass (g)		17754.1		17968.4	
Total pellets		228		257	
Food-niche breadth (H')		0.495		0.557	
Evenness (J')		0.519		0.584	
Geometric mean weight (g)		28.3		33	
Simple mean weight (g)		44.2		61.6	

^a Mass obtained from Jaksic et al. 1986.

^b Mass obtained from Martínez et al. 1998.

^c Mean mass of the two *Rattus* species inhabiting study areas (*R. rattus* and *R. norvegicus*).

^d Mean mass of all identified rodent prey.

ranking of prey species in the field (see Fig. 1). Only the southern big-eared mouse and the olivaceous field mouse appeared to have been consumed in a higher proportion than those estimated in the field (see Fig. 1).

Discussion

Barn owls and short-eared owls in our study area had very similar diets as they virtually preyed on the same rodent species in the same proportion. Convergence in hunting modes, activity time and body mass of owls, and prey abundance could be acting together. Both the barn and short-eared owl hunt individually, mainly by active searching and secondarily by sit-and-wait or perching (Jaksic & Carothers 1985). They are both very active flyers with cruising or quartering habits (Marti 1974, Jaksic & Carothers 1985). In this way, both owl species could have access to rodent prey with similar behaviour and microhabitat utilisation which should be mirrored in a high dietary similarity. However, some authors have suggested that the hunting modes are not steady predictors of food-niche metrics (e.g. Jaksic & Carothers 1985).

Even though barn and short-eared owls utilise similar habitats in our study area, hunting could have

taken place within defended territories (Korpimäki 1987, Holt & Leasure 1993). However, mean size of owls' territory may be particularly small in areas with very high small mammal abundance (Lockie 1955, Village 1987, Shaw 1995). Because the higher seasonal rodent abundance in agricultural areas of southern Chile occurs during late autumn and winter (R.A. Figueroa, unpubl. data), it is probable that barn and short-eared owls shared the same habitats, and therefore the same prey species, by reduction of hunting range when they concurrently searched for prey. Rodents may have provided an abundant food resource and thus made the level of exploitation competition insignificant (Jaksic 1985, Walk 1998), consequently leading to a high food-niche overlap (Lack 1946). Dietary similarities between barn owl and short-eared owl could also be due to the fact that their mean body masses are similar: 350-400 g and 400-500 g, respectively (Clark 1975, Marti 1992). This would guarantee access to prey species with similar body size. However, even though raptors' body mass has been demonstrated to be a good predictor of the size of their prey, it does not appear to influence diet breadth and overlap (Jaksic 1989).

Although differences in activity patterns between predators and prey generally could explain differential predation on rodent species, for the owl

species studied here, this does not appear to be the case. Barn owls are mostly nocturnal (Smith et al. 1974, Marti 1974), but occasionally hunt during day (e.g. Harte 1954, Haverschmidt 1970). Short-eared owls are also primarily nocturnal or crepuscular during winter (Clark 1975, Bosakowski 1989, Holt & Leasure 1993), but they may largely extend hunting activities to daylight hours (Clark 1975, Village 1987). In fact, on several occasions, we observed them searching for prey at midday and in the afternoon during winter. A similar daily activity pattern was observed in South Scotland by Village (1987). In this manner, it would be expected that the diet of both owl species did not overlap considerably, because they could access rodent prey with different daily cycles. However, most rodent prey species in southern Chile may extend their activity period largely into the day during winter (Muñoz et al. 1990), thus being available to owls both during the day and during the night. The diurnal activity of rodents during winter could have been accentuated by the shortening of daylight hours. Possibly, diurnal activity of short-eared owls avoided interference interactions with barn owls during night to a certain extent (Walk 1998, Leveau et al. 2004), but it could not be sufficient to cause marked differences in proportion of consumed rodent prey. Finally, the high homogeneity of the agricultural landscape in our study area would explain the high diet similarity among barn and short-eared owls from distinct localities, because this factor could determine relatively similar rodent prey assemblages on the most sites (Martínez et al. 1998, Bosè & Guidali 2001).

Our results suggest that barn and short-eared owls were opportunistic rather than selective predators in agricultural areas of the Osorno province, i.e. they appeared to prey more on the most abundant rodent prey species, the olivaceous field mouse and long-tailed pygmy rice rat (see Fig. 1). Various authors have claimed that barn owls show no food preferences, and that their diet is a reflection of the abundance of small mammals species within their hunting territories (Glue 1971, Bunn et al. 1982, Torres-Mura & Contreras 1989, Scheibler & Christoff 2004, Begall 2005). Similarly, short-eared owls also take whatever prey is most abundant or vulnerable to them (Martínez et al. 1998). However, in some studies, the proportion of consumed rodents by barn owls differed from those obtained by trapping in the field, suggesting prey selection (Perrins 1982, Bellocq 1998, Baxter & Matshili 2003). In addition, computer simulations have shown that barn owl pellets

could not truly mirror the proportion of prey in the field (Yom-Tov & Wool 1997). We recognise that our interpretation regarding prey selection could be erroneous, because most rodent trapping was not carried out concurrently with the collection of pellets. Moreover, results of live trapping are affected by sampled area, trap type and bait used, and could fail to capture all small mammal species present in the local community (Blem et al. 1993, Torre et al. 2004).

In addition to prey abundance, it is probable that activity level (Kaufman 1974) and microhabitat preference (Trejo & Guthmann 2003) could also explain the high proportion of long-tailed pygmy rice rat in the diets of both owls. This rat is a highly mobile species with a wider home range (0.032-0.480 ha) than the olivaceous field mouse (0.073-0.253 ha; Murúa et al. 1986) which would make it more vulnerable to predation. Moreover, the long-tailed pygmy rice rat in agricultural lands preferentially utilise dense shrubs mixed with pastures on fence borders (R.A. Figueroa unpubl. data), which increases the probability of encounters with owls, because fence posts are favourite perches (Martínez et al. 1998). In the case of the olivaceous field mouse, the occupation of more open habitat (e.g. abandoned pastures) could have increased its vulnerability (Dickman et al. 1991, Trejo & Guthmann 2003). Possibly, southern big-eared mouse was 'selected' because of its larger body size (51 g) relative to the remaining native rodents (≤ 30 g). We believe that higher GMWP for the short-eared owl was caused by the fact that it preyed on the largest rodent prey in our study sites, the Norwegian rat. Because it was not possible to identify most *Rattus* spp. remains from pellets at the species level, we did not know if Norwegian rats were eaten, and to what extent, by barn owls. Since GMWP may vary locally for the barn owl (Alvizatos & Goutner 1999, Pillado & Trejo 2000, Lekunze et al. 2001, Scheibler & Christoff 2004) and short-eared owl (Jaksic 1983, Cirignoli et al. 2001), our results may reflect the local body size distribution of the rodent community in agricultural areas of southern Chile.

Acknowledgements - we thank R. Murúa (Austral University of Chile) and J. Cerda (Agricultural and Livestock Service, Chile), who permitted access to the database of the rodent weights. A. Trejo, O. Skewes and R. González sent us some of the cited articles. This paper is a product of the Wildlife in Agricultural Landscape project financed by private funds. Most of the field work was financed by the grant FONDECYT 194-1256 from Chile's Fondo

Nacional de Investigación Científica y Tecnológica concedido a D.R. Martínez. R.A. Figueroa was supported by a Doctoral grant from Comisión Nacional de Investigación Científica y Tecnológica (CONICYT) and Escuela de Graduados, Facultad de Ciencias, Universidad Austral. Suggestions by Beatriz Arroyo helped to improve a first version of the manuscript.

References

- Alvizatos, H. & Goutner, V. 1999: Winter diet of the Barn Owl (*Tyto alba*) and Long-eared Owl (*Asio otus*) in northeastern Greece: a comparison. - *Journal of Raptor Research* 33: 160-163.
- Baxter, R.M. & Matshili, A.J. 2003: An analysis of Barn and Grass Owl pellets from Alice, Eastern Cape, South Africa. - *Ostrich* 74: 233-235.
- Begall, S. 2005: The relationship of foraging habitat to the diet of Barn Owls (*Tyto alba*) from central Chile. - *Journal of Raptor Research* 39: 97-101.
- Belloq, M.I. 1998: Prey selection by breeding and non-breeding Barn Owls in Argentina. - *Auk* 115: 224-229.
- Belloq, M.I. & Kravetz, F.O. 1994: Feeding strategy and predation of the Barn Owl (*Tyto alba*) and the Burrowing Owl (*Speotyto cunicularia*) on rodents species, sex, and size, in agrosystems of central Argentina. - *Ecología Austral* 4: 29-34.
- Blem, C.R., Blem, L.B., Felix, J.H. & Holt, D.W. 1993: Estimation of body mass of voles from crania in Short-eared owl pellets. - *American Midland Naturalist* 129: 281-286.
- Bosakowski, T. 1989: Observations on the evening departure and activity of wintering Short-eared Owls in New Jersey. - *Journal of Raptor Research* 23: 162-166.
- Bosè, M. & Guidali, F. 2001: Seasonal and geographic difference in the diet of the Barn Owl in an agroecosystem in northern Italy. - *Journal of Raptor Research* 35: 240-246.
- Bozinovic, F. & Medel, R. 1988: Body size, energetic and foraging mode of raptors in central Chile: an inference. - *Oecologia* 75: 456-458.
- Bunn, D.S., Warburton, A.B. & Wilson, R.D.S. 1982: The Barn Owl. - T. & A.D. Poyser, London, UK, 264 pp.
- Cirignoli, S., Podestá, D.H. & Pardiñas, U.F.J. 2001: Diet of the Short-eared Owl in northwestern Argentina. - *Journal of Raptor Research* 35: 68-69.
- Clark, R.J. 1975: A field study of the Short-eared Owl, *Asio flammeus* (Pontoppidan), in North America. - *Wildlife Monograph* 47: 1-67.
- di Castri, F. & Hajek, E. 1976: *Bioclimatología de Chile*. - Universidad Católica de Chile, Santiago, Chile, 129 pp.
- Dickman, C.R., Prevadec, M. & Lynam, A.J. 1991: Differential predation of size and sex classes of mice by the barn owl, *Tyto alba*. - *Oikos* 62: 67-76.
- Fowler, J. & Cohen, L. 1986: *Statistics for Ornithologist*. - British Trust for Ornithology, Norfolk, England, 150 pp.
- Fulk, G.W. 1976: Owl predation and rodent mortality: a case study. - *Mammalia* 40: 423-427.
- García, J.T. & Arroyo, B.E. 2005: Food-niche differentiation in sympatric Hen *Circus cyaneus* and Montagu's Harriers *Circus pygarcus*. - *Ibis* 147: 144-154.
- Gerstell, A.T. & Bednarz, J. 1999: Competition and patterns of resource use by two sympatric raptors. - *Condor* 101: 557-565.
- Glue, D.E. 1971: Avian predator pellet analysis and the mammalogist. - *Mammal Review* 21: 200-210.
- Harte, K. 1954: Barn Owl hunting by daylight. - *Wilson Bulletin* 66: 270.
- Haverschmidt, F. 1970: Barn Owls hunting by daylight in Surinam. - *Wilson Bulletin* 82: 101.
- Herrera, C.M. & Hiraldo, F. 1976: Food-niche and trophic relationships among European owls. - *Ornis Scandinavica* 7: 29-41.
- Holt, D.W. & Leasure, S.M. 1993: Short-eared Owl (*Asio flammeus*). - In: Poole, A. & Gill, F. (Eds.); *The Birds of North America*, No 62. Academy of Natural Science, Philadelphia, Pennsylvania, USA, and American Ornithological Union, Washington, D.C., USA, 24 pp.
- Holt, D.W., Lyon, L.J. & Hale, R. 1987: Techniques for differentiating pellets of Short-eared Owls and Northern Harriers. - *Condor* 89: 929-931.
- Jaksic, F.M. 1982: Inadequacy of activity time as a niche difference: the case of diurnal and nocturnal raptors. - *Oecologia* 52: 171-175.
- Jaksic, F.M. 1983: The trophic structure of sympatric assemblages of diurnal and nocturnal birds of prey. - *American Midland Naturalist* 109: 152-162.
- Jaksic, F.M. 1985: Toward raptor community ecology: behavior bases of assemblage structure. - *Journal of Raptor Research* 19: 107-112.
- Jaksic, F.M. 1989: Tamaño corporal de depredadores como predictor de atributos tróficos: el caso de las aves rapaces. - *Medio Ambiente (Chile)* 10: 23-26. (In Spanish).
- Jaksic, F.M. & Braker, H.E. 1983: Food-niche relationships and guild structure of diurnal birds of prey: competition versus opportunism. - *Canadian Journal of Zoology* 61: 2230-2241.
- Jaksic, F.M. & Carothers, J.H. 1985: Ecological, morphological, and bioenergetic correlates of hunting mode in hawks and owls. - *Ornis Scandinavica* 16: 165-172.
- Jaksic, F.M., Yañez, J.L. & Rau, J.R. 1986: Prey and trophic ecology of Great Horned Owls in western South America: an indication of latitudinal trends. - *Raptor Research* 20: 113-116.
- Kaufman, D.W. 1974: Differential predation on active and inactive prey by owls. - *Auk* 91: 172-173.
- Krebs, C.J. 1989: *Ecological methodology*. - Harper Collins Publisher, Inc., New York, 620 pp.

- Korpimäki, E. 1987: Dietary shifts, niche relationships and reproductive output of coexisting Kestrels and Long-eared Owls. - *Oecologia* 74: 277-285.
- Lack, D. 1946: Competition for food by birds of prey. - *Journal of Animal Ecology* 15: 123-129.
- Lekunze, L.M., Ezealor, A.U. & Aken'Ova, T. 2001: Prey groups in the pellets of the barn owl *Tyto alba* (Scopoli) in the Nigerian savanna. - *African Journal of Ecology* 39: 38-44.
- Leveau, L.M., Leveau, C.M. & Pardiñas, U.F.J. 2004: Trophic relationships between White-tailed Kites (*Elanus leucurus*) and Barn Owls (*Tyto alba*) in southern Buenos Aires province, Argentina. - *Journal of Raptor Research* 38: 178-181.
- Lockie, J.D. 1955: The breeding habits and food of Short-eared Owls after a vole plague. - *Bird Study* 2: 53-69.
- Marti, C. 1974: Feeding ecology of four sympatric owls. - *Condor* 76: 45-61.
- Marti, C. 1987: Raptor food habits studies. - In: Pendleton, B.A., Millsap, B.A., Cline, K.W. & Bird, D.M. (Eds.); *Raptor Management Techniques Manual* National Wildlife Federation, Washington D.C., pp. 67-79.
- Marti, C. 1992: Barn Owl (*Tyto alba*). - In: Poole, A., Stettenheim, P. & Gill, F. (Eds.); *The Birds of North America*, No 1. Academy of Natural Science, Philadelphia, Pennsylvania, USA, and American Ornithological Union, Washington D.C., USA, pp. 1-15.
- Marti, C.D. & Kochert, M.N. 1995: Are Red-tailed Hawks and Great Horned Owls diurnal-nocturnal counterparts? - *Wilson Bulletin* 107: 615-628.
- Martínez, D.R., Figueroa, R.A., Ocampo, C.L. & Jaksic, F.M. 1998: Food habits and hunting ranges of Short-eared Owls (*Asio flammeus*) in agricultural landscapes of southern Chile. - *Journal of Raptor Research* 32: 111-115.
- Muñoz, A., Murúa, R. & González, L.A. 1990: Nicho ecológico de la asociación de micromamíferos en un agroecosistema forestal de Chile central. - *Revista Chilena de Historia Natural* 63: 267-277. (In Spanish).
- Murúa, R., González, L.A. & Meserve, P.L. 1986: Population ecology of *Oryzomys longicaudatus* Philippii (Rodentia: Cricetidae) in southern Chile. - *Journal of Animal Ecology* 55: 281-293.
- Pearson, O. 1995: Annotated keys for identifying small mammals living in or near Nahuel Huapi National Park or Lanin National Park, southern Argentina. - *Mastozoología Neotropical* 2: 99-148.
- Perrins, M.R. 1982: Prey specificity of the Barn Owl in the Great Fish River valley of the Eastern Cape Province. - *South Africa Journal of Wildlife Research* 12: 14-25.
- Pillado, M.S. & Trejo, A. 2000: Diet of the Barn Owl (*Tyto alba tuidara*) in Northwestern Argentine Patagonia. - *Journal of Raptor Research* 34: 334-338.
- Rau, J.R., Martínez, D.R. & Yáñez, J. 1985: Dieta de la lechuza blanca, *Tyto alba* (Strigiformes, Strigidae) en el sur de Chile. - *Boletín del Museo Regional de la Araucanía* (Chile) 2: 134-135. (In Spanish).
- Rau, J.R., Villagra, M.C., Mora, M.L., Martínez, D.R. & Tillería, M.S. 1992: Food habits of the Short-eared Owl (*Asio flammeus*) in southern South America. - *Journal of Raptor Research* 26: 35-36.
- Reise, D. 1973: Clave para la determinación de los cráneos de marsupiales y roedores chilenos. - *Gayana (Zoolo-gía)* 27: 1-20. (In Spanish).
- Scheibler, D.R. & Christoff, A.U. 2004: Small mammals in the diet of Barn Owls (*Tyto alba*) in agroecosystems of southern Brazil. - *Ornitología Neotropical* 15: 65-70.
- Shaw, G. 1995: Habitat selection by Short-eared Owls *Asio flammeus* in young coniferous forests. - *Bird Study* 42: 158-164.
- Smith, D.G., Wilson, C.R. & Frost, H.H. 1974: History and ecology of a colony of Barn Owls in Utah. - *Condor* 76: 131-136.
- Steenhof, K. & Kochert, M.N. 1985: Dietary shifts of sympatric buteos during a prey decline. - *Oecologia* 66: 6-16.
- Torre, I., Arrizabalaga, A. & Flaquer, C. 2004: Three methods for assessing richness and composition of small mammal communities. - *Journal of Mammalogy* 85: 524-530.
- Torres-Mura, J.C. & Contreras, L.C. 1989: Ecología trófica de la lechuza blanca (*Tyto alba*) en los Andes de Chile Central. - *Studies on Neotropical Fauna and Environment* 24: 97-103. (In Spanish).
- Trejo, A. & Guthman, N. 2003: Owl selection on size and sex classes of rodents: activity and microhabitat use of prey. - *Journal of Mammalogy* 84: 652-658.
- Village, A. 1987: Numbers, territory-size and turnover of Short-eared Owls *Asio flammeus* in relation to vole abundance. - *Ornis Scandinavica* 18: 198-204.
- Walk, J.W. 1998: Winter roost sites of Northern Harriers and Short-eared Owls on Illinois grasslands. - *Journal of Raptor Research* 32: 116-119.
- Walker, R.S., Novaro, A.J., Perovic, P., Palacios, R., Donadio, E., Lucherini, M., Pía, M. & López, M.S. 2007: Diets of three species of andean carnivores in high-altitude deserts of Argentina. - *Journal of Mammalogy* 88: 519-525.
- Yom-Tov, Y. & Wool, D. 1997: Do the contents of Barn Owl pellets accurately represent the proportion of prey species in the field? - *Condor* 99: 972-976.
- Zar, J.H. 1996: *Biostatistical analysis*. 3rd edition. - Prentice Hall, Upper Saddle River, New Jersey, USA, 662 pp.